CONTENTS

The structure of an antigenic glycolipid (SL-IV) from Mycobacterium tuberculosis	
H.H. Baer (Ottawa, ON, Canada)	1
Synthesis and spectroscopic characterization of hydroxycinnamoylated methyl α -L-arabinofuranosyl- $(1 \rightarrow 2)$ - and $(1 \rightarrow 3)$ - β -D-xylopyranosides	
R.F. Helm and J. Ralph (Madison, WI, USA)	23
The crystal and molecular structure of the trisaccharide erlose as the monohydrate	
T. Taga, E. Inagaki, Y. Fujimori (Kyoto, Japan) and S. Nakamura (Okayama, Japan)	39
The X-ray crystal structure of N-(1-octyl)-D-talonamide and a consideration of its hydrogen-bonding scheme	
C. André, P. Luger, S. Svenson and JH. Fuhrhop (Berlin, Germany)	47
The application of NMR-pattern-recognition methods to the classification of reduced, peracety- lated oligosaccharide residues	
W.J. Goux and D.S. Weber (Richardson, TX, USA)	57
Structural characterization and rheological properties of an extracellular glucuronan produced by a <i>Rhizobium meliloti</i> M5N1 mutant strain	
A. Heyraud (Grenoble, France), J. Courtois (Amiens, France), L. Dantas, P. Colin-Morel (Grenoble, France) and B. Courtois (Amiens, France)	71
Contribution of the O-benzoyl groups of tetra-O-benzoyl- α,β -L-arabinopyranoses to the formation of 1,1-bis(benzamido)-1-deoxy-L-arabinitol	
S.R. Leicach, R.A. Cadenas and J.F. Sproviero (Buenos Aires, Argentina)	79
Fluorinated carbohydrates as potential plasma membrane modifiers. Synthesis of 3-deoxy-3-fluoro derivatives of 2-acetamido-2-deoxy-p-hexopyranoses	
M. Sharma, R.J. Bernacki, M.J. Hillman and W. Korytnyk (Buffalo, NY, USA)	85
The preparation and susceptibility to hydrolysis of novel O-galacturonoyl derivatives of carbohydrates	
J.A. Brown and S.C. Fry (Edinburgh, United Kingdom)	95
Synthesis, conformational analysis, and the glycosidic coupling reaction of substituted 2,7-di- oxabicyclo[4.1.0]heptanes: 1,2-anhydro-3,4-di- <i>O</i> -benzyl-β-L- and β-D-rhamnopyranoses	
Q. Chen, F. Kong and L. Cao (Beijing, People's Republic of China)	107
Syntheses and insulin-like activity of phosphorylated galactose derivatives	
HN. Caro, M. Martín-Lomas and M. Bernabé (Madrid, Spain)	119
Bile alcohol glucuronides: regioselective <i>O</i> -glucuronidation of 5β -cholestane- 3α , 7α , 12α , 25 -tetrol and 24 -nor- 5β -cholestane- 3α , 7α , 12α , 25 -tetrol	
B. Dayal, G. Salen, J. Padia, S. Shefer, G.S. Tint (Newark, NJ, USA), G. Sasso and T.H.	122
Williams (Nutley, NJ, USA)	133

Synthesis of 4-O- and 6-O-monoacryloyl derivatives of sucrose by selective hydrolysis of 4,6-O- (1-ethoxy-2-propenylidene)sucrose. Polymerization and copolymerization with styrene E. Fanton, C. Fayet, J. Gelas (Aubière, France), A. Deffieux, M. Fontanille and D. Jhurry	
(Talence, France)	143
Biosynthesis of curdlan from culture media containing ¹³ C-labeled glucose as the carbon source	
A. Kai, T. Ishino, T. Arashida (Tokyo, Japan), K. Hatanaka, T. Akaike (Yokohama,	
Japan), K. Matsuzaki (Ueda, Japan), Y. Kaneko and T. Mimura (Tokyo, Japan)	153
The synthesis of chemically modified disaccharide derivatives of the Shigella flexneri Y polysac- charide antigen	
FI. Auzanneau, H.R. Hanna and D.R. Bundle (Ottawa, ON, Canada)	161
A novel fructoglucan from the thermal polymerization of sucrose	
M. Manley-Harris and G.N. Richards (Missoula, MT, USA)	183
Anomerization and transglycosylation reactions of permethylated methyl D-glycopyranosides	
C.K. Lee, E.J. Kim and IS.H. Lee (Chuncheon, Korea)	197
Studies on the cell-wall mannan of the siphonous green algae, Codium latum	
S. Kaihou, T. Hayashi, O. Otsuru and M. Maeda (Saitama, Japan)	207
Phase diagrams of pectin-calcium systems: Influence of pH, ionic strength, and temperature on	
the gelation of pectins with different degrees of methylation	
C. Garnier, M.A.V. Axelos and JF. Thibault (Nantes, France)	219
Isolation and characterisation of the two main cell-wall types from guava (Psidium guajava L.)	
pulp O Manadia B Williams and I M Brillouet (Mantaellian France)	222
O. Marcelin, P. Williams and JM. Brillouet (Montpellier, France)	233
Studies on the regioselectivity of cellulose sulfation in an N ₂ O ₄ -N,N-dimethylformamide-cellulose system	
W. Wagenknecht, I. Nehls and B. Philipp (Teltow-Seehof, Germany)	245
Branching of amylose by the branching isoenzymes of maize endosperm	
Y. Takeda, HP. Guan and J. Preiss (East Lansing, MI, USA)	253
Structures of B90 (sugary) and W64A (normal) maize starches	
Y. Takeda and J. Preiss (East Lansing, MI, USA)	265
Characterization of the O-polysaccharide of Pasteurella haemolytica serotype A1	
W.B. Severn and J.C. Richards (Ottawa, ON, Canada)	277
Notes	
Standardized intermediates for oligosaccharide synthesis. A convenient preparation of partially protected derivatives of allyl $O-\beta$ -D-galactopyranosyl- $(1 \rightarrow 4)-\beta$ -D-glucopyranoside suitable for chain extension at position O-4'	
R.H. Youssef, B.A. Silwanis, R.I. El-Sokkary (Alexandria, Egypt), A.S. Nematalla and	
M.A. Nashed (Alameda, CA, USA)	287
Synthesis and conformation of 1,2-anhydro-3,4-6-tri-O-benzyl-β-D-talopyranose	
I Liu F Kong and I Cao (Reijing People's Penublic of China)	205

_						
Contents	/ Carbohydr.	Res.	240	(1993)	C23-	-C25

	e	ъ,	~	e
-)			.,	-

Preparation of 2-amino-2-deoxy-3,4,5,6-tetra-O-methyl-p-gluconic acid hydrochloride, an intermediate in the preparation of polypeptide-type polyamides	
M. de Gracia García Martín, M.V. de Paz Báñez and J.A. Galbis Pérez (Sevilla, Spain) .	301
Synthesis of low-toxicity, 5-deoxy-5-fluoro and 5-deoxy-5,5-difluoro derivatives of arbekacin and its analogs, and study of structure-toxicity relationships	
T. Tsuchiya, T. Shitara, S. Umezawa (Kawasaki, Japan), T. Takeuchi, M. Hamada (Tokyo,	
Japan), N. Tomono and E. Umemura (Yokohama, Japan)	307
Synthesis of cyclophosphamide analogs from aminotrideoxy sugars	
C. Monneret, R. Gagnet and JC. Florent (Paris, France)	313
Author index	C1
Subject index	C3
Cumulative Author index (Vols. 221–240)	C7
Contents	C23

